



COMPREHENSIVE

UROLOGIC CARE

RICHARD B. TROY MD, FACS
NING Z. WU, MD, PHD
TAMRA E. LEWIS, MD, FACS
BRIAN M. KEUER, MD
CHRISTOPHER M. LODOWSKY, MD
ANDREA SCHRAGE, PA-C
SAMANTHA L. POHLMAN, PA-C
VICKI SZORC, PA-C
TAYLOR JOHNSON, FNP
DAVID DEGRASSI, FNP-BC

Extracorporeal Shockwave Lithotripsy (ESWL) Postoperative Instructions

DIET:

You may resume a normal diet as soon as you feel ready after your procedure. It is common for 24 hours after anesthesia to avoid heavy meals and start slowly with clear liquids. Eat plenty of fiber-rich foods for the first 1-2 weeks. Alcoholic beverages are best avoided in the first 24 hours and while taking any narcotic pain medications. It is important to drink plenty of water (6-8 glasses daily) after your ESWL.

ACTIVITY:

It is important to avoid strenuous activity and/or exercise for the first week after your procedure. This includes avoiding golfing, tennis, and jogging, working out/stretching activities, lawn work such as mowing, raking, shoveling, sexual activity, and any lifting more than 15 lbs. Discuss any specific work-related concerns you have with your urologist.

MEDICATIONS:

Use Tylenol for pain as directed. Ibuprofen can be used starting day 3.

It is important that if you take a daily blood-thinning/anti-platelet medication (Coumadin/warfarin, aspirin, Advil/Motrin, Lovenox, Pradaxa, Eliquis, Xeralto, Plavix, etc.) that you only restart this as instructed by your physicians.

BOWELS:

It is important to avoid constipation as you recover from your ESWL. Drink plenty of fluids (especially water). Narcotic pain medication can be constipating and it is important to take an over-the-counter stool softener/laxative to help prevent constipation (Miralax, milk of magnesia, Colace, etc.).

22285 N. Pepper Road, Suite 201
Lake Barrington, IL 60010-2540

1600 N. Randall Road, Suite 201
Elgin, IL 60123-7805

1585 N Barrington Road, Suite 206
Hoffman Estates, IL 60169

847.382.5080 T | 847.382.0923 F

www.compurocare.com

EXPECTED SYMPTOMS:

It is common to see blood in the urine for several days after your procedure. You may see sediment or stone fragments pass in the urine after your procedure. If you were given a urinary strainer, strain your urine and save any fragments. In some cases a ureteral stent is placed at the time of your ESWL-this can cause urinary urgency/frequency, burning with urination, bladder spasms, and pain in kidney with voiding. Some redness or bruising at the site where the machine was focused to break-up your stones can be normal.

Contact us for severe pain and/or burning with urination, fevers and chills, severe bleeding with passage of large blood clots, or inability to pass urine.

FOLLOWUP APPOINTMENT:

Our office will contact you to schedule your follow-up appointment. This will typically be in 1-2 weeks to check a KUB X-ray and possibly remove your stent.